

Effectiveness of the Basic Infection Control Skills License (BICSL) Training Program on Nurses' Knowledge, Attitudes, and Practices in Makkah, Saudi Arabia



Azizah Banafi^{1,2,*} and Noor Mastura Mohd Muiar^{2,*} 

¹Medical Affairs Unit, Executive Administration of Primary Healthcare Centers, Makkah, Saudi Arabia

²Department of Community Health, Cancer Research and Specialist Centre, Universiti Sains Malaysia (USM), 13200 Kepala Batas, Penang, Malaysia

Abstract:

Introduction: Healthcare-associated infections (HCAs) remain a major global health concern. In Saudi Arabia, the Basic Infection Control Skills License (BICSL) program was introduced to standardize infection prevention and control (IPC) training among healthcare workers. However, evidence on its effectiveness among nurses remains limited.

Methods: A multicenter, quasi-experimental, one-group pre-post study was conducted among 360 nurses from 5 public hospitals in Makkah, Saudi Arabia, during the 2024 BICSL implementation cycle. All participants received the same 3-day, in-person BICSL intervention; no random allocation or concurrent comparison group was used. A validated self-administered questionnaire assessed knowledge, attitude, and self-reported practice before and immediately after training. Paired t-tests, Wilcoxon signed-rank tests, effect sizes, reliability analyses, baseline-adjusted models, change-score models, and sensitivity analyses were conducted.

Results: Knowledge scores increased from 8.54 ± 2.03 to 10.71 ± 1.79 , attitude scores from 2.30 ± 0.77 to 2.70 ± 0.51 , and practice scores from 4.48 ± 1.91 to 5.90 ± 1.85 (all $p < 0.001$). Wilcoxon tests confirmed the pre-post differences. Cohen's d values were 0.72 for knowledge, 0.43 for attitude, and 0.55 for practice. In adjusted analyses, improvement in knowledge remained significantly associated with post-training practice and practice improvement.

Discussion: The findings suggest that structured BICSL training may strengthen nurses' IPC-related knowledge and self-reported practice. However, interpretation should remain cautious because the one-group pre-post design cannot exclude testing effects, seasonal influences, concurrent IPC activities, or social desirability bias.

Conclusion: BICSL training was associated with improved knowledge, attitudes, and practices (KAP) scores among nurses in hospitals in Makkah. Future evaluations should include control groups, longer follow-up, observational audits, and IPC outcome indicators.

Keywords: Infection prevention and control, Standard precautions, Basic infection control skills license (BICSL), Knowledge-attitude-practice (KAP), Nurses, Saudi Arabia, Healthcare-associated infections, Training effectiveness.

© 2026 The Author(s). Published by Bentham Open.

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

*Address correspondence to these authors at the Department of Community Health, Cancer Research and Specialist Centre, Universiti Sains Malaysia (USM), 13200 Kepala Batas, Penang, Malaysia and Medical Affairs Unit, Executive Administration of Primary Healthcare Centers, Makkah, Saudi Arabia; E-mails: masturamuiar@usm.my and azizahbanafi2025@gmail.com

Cite as: Banafi A, Muiar N. Effectiveness of the Basic Infection Control Skills License (BICSL) Training Program on Nurses' Knowledge, Attitudes, and Practices in Makkah, Saudi Arabia. Open Nurs J, 2026; 20: e18744346494646. <http://dx.doi.org/10.2174/0118744346494646260824073215>



Received: April 16, 2026
Revised: June 22, 2026
Accepted: July 07, 2026
Published: August 28, 2026



Send Orders for Reprints to
reprints@benthamscience.net

1. INTRODUCTION

Healthcare-associated infections (HCAIs) are a major cause of morbidity, mortality, and increased healthcare costs worldwide. HCAI burden varies across regions and remains a continuing challenge in both high-income and emerging healthcare systems [1, 2]. In Saudi Arabia, hospital infections remain a recognized public health issue, and previous studies have reported gaps in knowledge of hand hygiene and adherence to infection control among healthcare workers [3-6].

Standard precautions are central to infection prevention and control (IPC). They include hand hygiene, appropriate use of personal protective equipment (PPE), sharps safety, environmental cleaning, injection safety, waste disposal, and precautions for transmission-based risks [7-9]. Nurses are frontline healthcare workers and are therefore central to IPC implementation in routine clinical care [10, 11].

The Kingdom of Saudi Arabia has implemented several strategies to strengthen IPC, including the Basic Infection Control Skills License (BICSL) program. BICSL was established to provide standardized training in essential IPC competencies across healthcare settings [12]. Although several Saudi studies have examined IPC knowledge, attitudes, and practices (KAP) among healthcare workers [13-17], limited evidence exists on whether BICSL training itself is associated with measurable changes in nurses' KAP.

This study evaluated changes in nurses' knowledge, attitudes, and self-reported practices regarding standard precautions after BICSL training in Makkah, Saudi Arabia. The study was guided by the KAP framework, recognizing that knowledge may shape attitudes and that both may influence practice, while also acknowledging that practice change is affected by organizational and contextual factors such as workload, resources, and institutional IPC culture [6, 11, 18, 19].

1.1. RESEARCH QUESTIONS

The following research questions guided this study; no formal directional hypotheses were prespecified:

- [1] To what extent do nurses' knowledge, attitudes, and self-reported practices regarding standard precautions change after completing the BICSL training program?
- [2] Are the observed pre-post changes in knowledge, attitude, and practice scores statistically significant when assessed using both parametric and non-parametric methods?
- [3] Which demographic, professional, and institutional factors are associated with post-training practice scores and improvement in practice after adjusting for baseline practice?
- [4] How should the observed improvements be interpreted considering the quasi-experimental one-group pre-post design and the absence of a control group?

2. MATERIALS AND METHODS

2.1. Research Design and Study Setting

This multicenter, quasi-experimental, one-group pre-post study was conducted in 5 Ministry of Health public hospitals in Makkah, Saudi Arabia: Al Noor Specialist Hospital, Hira General Hospital, King Abdulaziz Hospital, King Faisal Hospital, and Maternity and Children Hospital. The study was implemented during the 2024 Hajj-related BICSL training period, when hospitals in Makkah experienced increased patient flow and heightened IPC preparedness requirements. All participants received the same intervention; there was no random allocation or concurrent comparison group. The study is reported in accordance with the TREND guideline for nonrandomized evaluations.

Because the study did not include randomization or a concurrent control group, the design assessed within-participant changes after training rather than establishing definitive causal effects. Potential threats to causal inference, including testing effects, maturation, secular trends, and concurrent IPC initiatives, were considered when interpreting the findings.

2.2. Target Population, Sampling, and Participants

The target population comprised registered nurses employed in the 5 participating hospitals who had not completed BICSL training during the current implementation cycle. Healthcare workers other than nurses, nurses who did not consent, and nurses without paired pre- and post-training data were ineligible for the paired analysis. The sampling frame included approximately 3,500 nurses in Makkah, and systematic random sampling was applied to the nursing staff lists, selecting approximately every 10th nurse. The minimum required sample size was 347; the calculation parameters were not available in the study documentation.

The study records documented 360 nurses who completed the pre-training questionnaire, the three-day BICSL training, and the immediate post-training questionnaire and who were included in the final paired analysis. The numbers initially approached or assessed for eligibility, excluded or declining before enrollment, and lost during follow-up were not separately documented. Protocol deviations were also not separately documented (Fig. 1).

2.3. BICSL Training Intervention

The BICSL intervention was delivered in person over 3 consecutive days by certified BICSL trainers from the participating hospitals' infection control departments. The standardized curriculum covered 7 core components: hand hygiene, personal protective equipment, biological spill management, needle-stick injury management, transmission-based precautions, respirator fit testing, and powered air-purifying respirators. Each module combined theoretical instruction, trainer demonstration, and participant return demonstration. Training was delivered in Arabic and English as needed. Because all participants

received the educational intervention, neither participants nor trainers were blinded. Although the curriculum and procedures were standardized across hospitals, formal independent fidelity auditing of individual sessions was not conducted. Session group size was not recorded, and no incentives or other adherence-promoting activities were documented.

2.4. Data Collection and Measurement Instrument

Data were collected using a self-administered structured questionnaire adapted from previously used KAP instruments and aligned with CDC and WHO IPC guidance [6, 20, 21]. The questionnaire captured demographic and professional characteristics and assessed 3 co-primary outcome domains: knowledge, attitude, and

practice. A pilot validity check was conducted with 30 nurses before the main study; these pilot observations were not included in the final analytic sample.

The knowledge scale included 13 dichotomous items scored 1 for a correct response and 0 for an incorrect or “do not know” response, giving a score range of 0-13. The attitude scale included three dichotomous items, scored 1 for a positive/correct response and 0 otherwise, giving a score range of 0-3. The practice scale included 9 self-reported items. For the primary analysis, “always” was coded as 1 to represent strict adherence, while “often,” “sometimes,” “seldom,” and “never” were coded as 0, yielding a score range of 0-9. This dichotomization was selected to identify consistent IPC compliance, but it also reduces ordinal information, which is noted as a limitation.




Fig. (1). Participant flow through recruitment, BICSL training, and paired analysis.

The pre-test was administered before BICSL training. The post-test was administered immediately after completion of the 3-day training within the same implementation cycle.

2.5. Statistical Analysis

Descriptive statistics summarized participant characteristics and KAP scores. The individual nurse was the unit of analysis. Paired t-tests were used to compare pre- and post-training KAP scores. Because the domain scores were based on dichotomous items and might not be normally distributed, Wilcoxon signed-rank tests were also reported. Domain-level effect sizes were calculated using Cohen's *dz*, and ceiling effects were assessed by reporting the proportion of participants who achieved the maximum score. No missing values were imputed; analyses were restricted to complete paired records. Data were analyzed using IBM SPSS Statistics version 31.0.2.0.

Internal consistency was assessed separately for each KAP subscale at pre-test and post-test using KR-20/Cronbach's alpha and average inter-item correlations. Because the attitude scale contained only 3 items and all domains used dichotomous scoring, reliability estimates were interpreted cautiously.

To address concerns about post-test-only regression, the revised analysis used baseline-adjusted and change-score models. The baseline-adjusted model estimated the post-training practice score, adjusting for baseline practice, changes in knowledge and attitude, gender, nationality, qualification, department group, and hospital fixed effects. The change-score model estimated change in practice score and included baseline practice, knowledge, and attitude, changes in knowledge and attitude, and the same covariates. HC3 robust standard errors were used. Continuous predictors were mean-centered before creating interaction terms. A binomial GLM sensitivity analysis modeled the number of compliant practice items out of 9. Variance inflation factors (VIFs) were examined;

age was evaluated but not retained in the final hospital fixed-effect models because diagnostics indicated instability when age and hospital indicators were included together. Baseline practice was retained in all adjusted models. Holm-Bonferroni correction was applied to exploratory pairwise subgroup comparisons.

2.6. Ethics

The study was approved by the Jawatankuasa Etika Penyelidikan Manusia Universiti Sains Malaysia (JEPeM-USM; protocol code USM/JEPeM/PP/23020204) and by the General Directorate of Research & Studies, Ministry of Health, Kingdom of Saudi Arabia (decision approval number 1122-836 076-K-02-H). Data were anonymized before analysis.

3. RESULTS

3.1. Participant Characteristics

Participant flow is presented in Fig. (1). A total of 360 nurses with complete paired pre- and post-training data were included. The mean age was 39.93 ± 11.00 years (range 22-59 years). Most participants were male (55.00%), Saudi (53.06%), staff nurses (42.78%), and employed in ICU departments (74.17%). Participant characteristics are presented in Table 1.

3.2. Pre- and Post-Training Kap Changes

KAP scores increased significantly across all domains after BICSL training (Table 2). Knowledge increased from 8.54 ± 2.03 to 10.71 ± 1.79 , attitude from 2.30 ± 0.77 to 2.70 ± 0.51 , and practice from 4.48 ± 1.91 to 5.90 ± 1.85 . Paired t-tests and Wilcoxon signed-rank tests were statistically significant for all domains ($p < 0.001$). Cohen's *dz* indicated a moderate-to-large effect for knowledge (0.72), a small-to-moderate effect for attitude (0.43), and a moderate effect for practice (0.55). Ceiling effects were most evident for attitude, for which 72.5% of participants achieved the maximum post-training score.

Table 1. Sociodemographic and professional characteristics of participants (n = 360).

Characteristic	Category	n	% or Value
Age, years	Mean $\pm$ SD (range)	-	39.93 ± 11.00 (22-59)
Hospital	Hira General Hospital	87	24.17
Hospital	Maternity and Children Hospital	76	21.11
Hospital	King Faisal Hospital	74	20.56
Hospital	Al Noor Specialist Hospital	63	17.50
Hospital	King Abdulaziz Hospital	60	16.67
Gender	Male	198	55.00
Gender	Female	162	45.00
Nationality	Saudi	191	53.06
Nationality	Non-Saudi	169	46.94
Qualification	Staff Nurse	154	42.78
Qualification	Senior Nurse	110	30.56
Qualification	Registered Nurse	96	26.67
Department group	ICU	267	74.17
Department group	Others	93	25.83

Table 2. Pre-post changes in KAP scores with non-parametric confirmation, effect sizes, and ceiling effects.

Domain	Max Score	Pre-training Mean $\pm$ SD	Post-training Mean $\pm$ SD	Mean Difference	95% CI	t(df = 359)	t-test p-value	Wilcoxon p-value	Cohen's dz	Pre at Maximum n (%)	Post at Maximum n (%)
Knowledge	13	8.54 $\pm$ 2.03	10.71 $\pm$ 1.79	2.17	1.86, 2.48	13.68	<0.001	<0.001	0.72	8 (2.2)	62 (17.2)
Attitude	3	2.30 $\pm$ 0.77	2.70 $\pm$ 0.51	0.40	0.30, 0.50	8.24	<0.001	<0.001	0.43	169 (46.9)	261 (72.5)
Practice	9	4.48 $\pm$ 1.91	5.90 $\pm$ 1.85	1.41	1.15, 1.68	10.45	<0.001	<0.001	0.55	2 (0.6)	38 (10.6)

Note: The mean difference was calculated as the post-training score minus the pre-training score. Cohen's dz represents the paired-sample standardized mean difference. The proportion of participants achieving the maximum score was reported to assess possible ceiling effects.

3.3. Reliability of KAP Subscales

Internal consistency was low, especially for the three-item attitude domain (Table 3). Accordingly, the KAP domain totals should be interpreted as pragmatic summary scores of heterogeneous IPC competencies rather than as highly internally consistent psychometric scales.

3.4. Baseline-adjusted and Change-score Models

In the baseline-adjusted model (Table 4), change in knowledge was significantly associated with a higher post-

training practice score ($\beta = 0.105$, $p = 0.001$). In contrast, change in attitude was positive but not statistically significant ($\beta = 0.150$, $p = 0.154$). Staff nurses had lower adjusted post-training practice scores than senior nurses ($\beta = -1.552$, $p < 0.001$). ICU nurses had higher adjusted practice scores than nurses in other departments ($\beta = 0.583$, $p = 0.007$). The model R^2 was 0.251, and the adjusted R^2 was 0.225.

The model used OLS with HC3 robust standard errors and hospital fixed effects. Reference groups were female, non-Saudi, senior nurse, non-ICU/other departments, and Hira General Hospital.

Table 3. Internal consistency of KAP subscales at pre-test and post-test.

Domain	Time Point	No. Items	Score Range	Mean $\pm$ SD	KR-20 / alpha	Avg Inter-item Correlation
Knowledge	Pre	13	0-13	8.54 $\pm$ 2.03	0.323	0.036
Knowledge	Post	13	0-13	10.71 $\pm$ 1.79	0.441	0.057
Attitude	Pre	3	0-3	2.30 $\pm$ 0.77	0.143	0.052
Attitude	Post	3	0-3	2.70 $\pm$ 0.51	-0.051	-0.017
Practice	Pre	9	0-9	4.48 $\pm$ 1.91	0.434	0.079
Practice	Post	9	0-9	5.90 $\pm$ 1.85	0.458	0.086

Note: KR-20/Cronbach's alpha values should be interpreted cautiously because the KAP domains contain heterogeneous IPC items and the attitude domain includes only 3 dichotomous items. The low post-training attitude reliability likely reflects limited item variance and a ceiling effect rather than absence of training response.

Table 4. Baseline-adjusted model for post-training practice score.

Variable	Beta	Robust SE	95% CI	p-value
Intercept	6.69	0.407	5.89, 7.49	<0.001
Male vs. Female	0.074	0.179	-0.28, 0.42	0.680
Saudi vs. Non-Saudi	-0.106	0.175	-0.45, 0.24	0.543
Staff Nurse vs. Senior Nurse	-1.552	0.218	-1.98, -1.13	<0.001
Registered Nurse vs. Senior Nurse	-0.463	0.243	-0.94, 0.01	0.057
ICU vs. Others	0.583	0.214	0.16, 1.00	0.007
Al Noor Specialist vs. Hira General	-1.245	0.309	-1.85, -0.64	<0.001
King Abdulaziz vs. Hira General	-0.724	0.284	-1.28, -0.17	0.011
King Faisal vs. Hira General	-0.689	0.252	-1.18, -0.19	0.006
Maternity & Children vs. Hira General	-0.607	0.249	-1.10, -0.12	0.015
Baseline practice score	-0.022	0.05	-0.12, 0.08	0.651
Change in knowledge score	0.105	0.032	0.04, 0.17	0.001
Change in attitude score	0.15	0.105	-0.06, 0.36	0.154

Note: Outcome: post-training practice score (0-9).

In the change-score model (Table 5), improvement in knowledge was significantly associated with improvement in practice ($\beta = 0.181$, $p < 0.001$). Improvement in attitude was positive but not statistically significant ($\beta = 0.280$, $p = 0.114$). Lower baseline practice was strongly associated with greater practice improvement ($\beta = -1.018$, $p < 0.001$), consistent with greater room for improvement among participants with lower initial scores. The model R^2 was 0.615 and adjusted R^2 was 0.599.

3.5. Sensitivity and Multiple-comparison Analyses

The binomial GLM sensitivity analysis treated practice as the number of compliant “always” responses out of 9 (Table 6). Results were consistent with the OLS models:

improvement in knowledge remained significantly associated with compliant practice responses ($OR = 1.055$, $p < 0.001$), whereas change in attitude was not statistically significant after adjustment ($OR = 1.081$, $p = 0.139$).

After Holm-Bonferroni correction, most exploratory subgroup differences were no longer statistically significant. For change-score subgroup comparisons, only the difference in attitude change between senior nurses and staff nurses remained statistically significant after correction (mean difference = -0.353 , Holm-adjusted $p = 0.041$). Therefore, all subgroup findings are interpreted as exploratory (Table 7).

Table 5. Change-score model for improvement in practice score.

Variable	Beta	Robust SE	95% CI	p-value
Intercept	4.831	1.029	2.81, 6.85	<0.001
Male vs. Female	0.074	0.179	-0.28, 0.43	0.681
Saudi vs. Non-Saudi	-0.116	0.174	-0.46, 0.23	0.506
Staff Nurse vs. Senior Nurse	-1.511	0.217	-1.94, -1.09	<0.001
Registered Nurse vs. Senior Nurse	-0.45	0.241	-0.92, 0.02	0.062
ICU vs. Others	0.545	0.216	0.12, 0.97	0.012
Al Noor Specialist vs. Hira General	-1.081	0.329	-1.73, -0.44	0.001
King Abdulaziz vs. Hira General	-0.615	0.286	-1.17, -0.05	0.032
King Faisal vs. Hira General	-0.607	0.256	-1.11, -0.11	0.018
Maternity & Children vs. Hira General	-0.551	0.253	-1.05, -0.05	0.030
Baseline practice score	-1.018	0.05	-1.12, -0.92	<0.001
Baseline knowledge score	0.136	0.069	0.00, 0.27	0.048
Baseline attitude score	0.175	0.219	-0.25, 0.60	0.425
Change in knowledge score	0.181	0.052	0.08, 0.28	<0.001
Change in attitude score	0.28	0.177	-0.07, 0.63	0.114

Note: Outcome: change in practice score (post-training minus pre-training). The model used OLS with HC3 robust standard errors and hospital fixed effects.

Table 6. Binomial GLM sensitivity analysis for bounded practice outcome.

Variable	Odds Ratio	95% CI Lower	95% CI Upper	p-value
Intercept	2.978	1.978	4.484	<0.001
Male vs. Female	1.04	0.872	1.24	0.660
Saudi vs. Non-Saudi	0.94	0.791	1.118	0.486
Staff Nurse vs. Senior Nurse	0.453	0.362	0.566	<0.001
Registered Nurse vs. Senior Nurse	0.771	0.595	1	0.050
ICU vs. Others	1.349	1.1	1.655	0.004
Al Noor Specialist vs. Hira General	0.525	0.388	0.711	<0.001
King Abdulaziz vs. Hira General	0.671	0.502	0.898	0.007
King Faisal vs. Hira General	0.687	0.527	0.895	0.005
Maternity & Children vs. Hira General	0.712	0.547	0.927	0.012
Baseline practice score	0.991	0.944	1.041	0.727
Change in knowledge score	1.055	1.023	1.089	<0.001
Change in attitude score	1.081	0.975	1.199	0.139

Note: Outcome: number of practice items rated “always” out of 9. The model used a logit link and HC3-robust standard errors.

Table 7. Summary of Holm-Bonferroni multiplicity correction.

Analysis set	Number of Pairwise Tests per KAP Domain	Findings Remaining Significant after Holm Correction
Change-score subgroup comparisons	16 per domain	Only attitude change differed significantly between senior nurses and staff nurses; mean difference = -0.353, Holm-adjusted $p = 0.041$.
Baseline subgroup comparisons	16 per domain	Baseline attitudes differed between senior nurses and staff nurses; baseline practices differed between senior nurses and staff nurses, and between staff nurses and registered nurses.

4. DISCUSSION

This study found statistically significant improvements in nurses' knowledge, attitudes, and self-reported practices after completion of the BICSL training program. The magnitude of improvement was greatest for knowledge, followed by practice and attitude. The findings are consistent with the KAP framework and recent IPC literature, which suggest that knowledge acquisition, skills reinforcement, and supportive clinical environments can contribute to better IPC practice [11, 18, 19].

The improvement in knowledge is consistent with previous IPC training studies showing that structured educational interventions can improve nurses' understanding of standard precautions, hand hygiene, personal protective equipment, and sharps safety [10, 19, 22-24]. In the present study, BICSL combined theoretical instruction with demonstration and return-demonstration. This practical structure may explain why gains in knowledge were accompanied by measurable improvements in self-reported practice. The moderate baseline knowledge score also allowed room for observable post-training improvement.

Attitude improved significantly, but the standardized effect size was smaller than those observed for knowledge and practice. This may be explained by a ceiling effect, because many participants already had favorable attitudes before training and the post-training attitude score approached the maximum possible value. This finding differs from studies in which lower baseline attitude scores allowed larger post-training gains. In this context, BICSL may be more effective at reinforcing existing positive attitudes and translating them into practice than at producing large, measurable shifts in attitude scores.

Practice scores improved after training, consistent with previous studies reporting better IPC compliance following educational interventions [11, 22-24]. However, this finding must be interpreted cautiously because practice was self-reported and dichotomized using a strict definition, with only "always" coded as compliant. Differences between the present findings and studies based on direct observation may reflect social desirability bias, differences in clinical workload, and variation in institutional IPC monitoring systems [11].

The adjusted analyses suggest that improvement in knowledge was associated with higher post-training practice scores and greater improvement in practice. In contrast, improvement in attitude was positive but not statistically significant after adjustment. This may indicate that practice change after BICSL depends strongly on

procedural knowledge and repeated competency demonstration, whereas favorable attitudes may be necessary but not sufficient for sustained behavioral change. Organizational conditions, such as workload, managerial support, availability of PPE, and other IPC resources, may influence whether favorable attitudes are translated into consistent practice [8, 18].

The higher adjusted practice scores among ICU nurses may reflect greater routine exposure to IPC procedures, higher perceived infection risk, and more frequent reinforcement of IPC policies in critical care settings. Conversely, differences by qualification and hospital site may reflect variation in staffing patterns, supervision, resource availability, local IPC culture, and opportunities for supervised practice [2, 11, 18, 25]. These findings indicate that BICSL effectiveness may depend not only on the training curriculum but also on the clinical and organizational environment in which nurses apply IPC practices.

5. LIMITATIONS

This study has several limitations. First, it used a quasi-experimental, one-group pre-post design without randomization or a concurrent control group. Therefore, although KAP scores improved after BICSL training, the changes cannot be attributed exclusively to the training program. Testing effects, repeated exposure to the questionnaire, maturation, secular trends, seasonal workload changes, Hajj-related preparedness activities, and concurrent institutional IPC initiatives may have contributed to the observed improvements.

Second, practice was measured using self-reported questionnaire items rather than direct observation or audit-based measures, which may have introduced social desirability bias. In addition, practice responses were dichotomized, with "always" coded as compliant and all other responses coded as non-compliant. Although this approach reflects strict adherence, it may have reduced measurement sensitivity by collapsing ordinal response information.

Third, the attitude scale contained only 3 items, and subscale reliability was low, particularly for the attitude domain. Therefore, the attitude findings should be interpreted cautiously. Future studies should use expanded and psychometrically validated KAP instruments with stronger construct validity.

Fourth, the study was conducted in 5 public hospitals in Makkah, and ICU nurses were overrepresented in the sample. Therefore, the findings may not be fully generalizable to non-ICU departments, primary healthcare

centers, private hospitals, or other regions of Saudi Arabia.

Fifth, the post-training assessment was conducted within the same implementation cycle and measured short-term change. The study did not assess long-term retention of knowledge, attitudes, or practices. Future studies should include follow-up assessments at 3, 6, and 12 months.

Finally, this study did not include direct patient-safety or IPC outcome indicators, such as hand hygiene audit data, PPE compliance audits, needle-stick injury reports, or healthcare-associated infection rates. Training-related adverse events or unintended effects were not systematically collected. Future BICSL evaluations should triangulate self-reported KAP outcomes with observational audits and institutional IPC surveillance data.

CONCLUSION

Structured BICSL training was associated with improved knowledge, attitude, and self-reported practice scores among nurses in Makkah public hospitals. These findings support BICSL as a practical strategy for building IPC competencies, but the one-group design requires cautious interpretation. Future evaluations should include control groups, longer follow-up, observational audits, and IPC outcome indicators to confirm sustainability and patient-safety relevance.

AUTHORS' CONTRIBUTIONS

The authors' contributions were as follows: A.A.B. and N.M.M.M.: Study conception and design; A.A.B.: Data collection; A.A.B., N.M.M.M., and R.H.: Analysis and interpretation of results; A.A.B.: Initial manuscript preparation; and N.M.M.M. and R.H.: Critical revision of the manuscript. All authors reviewed the results and approved the final version of the manuscript.

LIST OF ABBREVIATIONS

BICSL	= Basic Infection Control Skills License
CBAHI	= Saudi Central Board for Accreditation of Healthcare Institutions
CDC	= Centers for Disease Control and Prevention
HAI/HCAI	= Healthcare-Associated Infection
HCW	= Healthcare Worker
ICU	= Intensive Care Unit
IPC	= Infection Prevention and Control
KAP	= Knowledge, Attitude, and Practice
KSA	= Kingdom of Saudi Arabia
MOH	= Ministry of Health
NSI	= Needle-Stick Injury
PAPR	= Powered Air-Purifying Respirator
PPE	= Personal Protective Equipment
SP	= Standard Precautions

WHO = World Health Organization

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study was approved by the Jawatankuasa Etika Penyelidikan Manusia Universiti Sains Malaysia (JEPeM-USM; protocol code USM/JEPeM/PP/23020204) and by the General Directorate of Research & Studies, Ministry of Health, Kingdom of Saudi Arabia (decision approval number 1122-836 076-K-02-H).

HUMAN AND ANIMAL RIGHTS

All human research procedures followed were in accordance with the ethical standards of the committee responsible for human experimentation (institutional and national), and with the Helsinki Declaration of 1975, as revised in 2013.

CONSENT FOR PUBLICATION

Participants provided consent before participation.

STANDARDS OF REPORTING

The study was reported in accordance with the TREND guidelines.

AVAILABILITY OF DATA AND MATERIALS

The data supporting the analysis are available from the authors upon reasonable request and upon approval from the Ministry of Health, Saudi Arabia.

FUNDING

None.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The authors thank the participating hospitals, nursing departments, BICSL trainers, and all nurses who completed the questionnaires.

REFERENCES

- [1] Haque M, Sartelli M, McKimm J, Abu Bakar MB. Health care-associated infections - an overview. *Infect Drug Resist* 2018; 11: 2321-33. <http://dx.doi.org/10.2147/IDR.S177247> PMID: 30532565
- [2] World Health Organization. Global report on infection prevention and control 2024. 2024. Available from :<https://www.who.int/citations/i/item/9789240103986>
- [3] Memish ZA. Infection control in Saudi Arabia: Meeting the challenge. *Am J Infect Control* 2002; 30(1): 57-65. <http://dx.doi.org/10.1067/mic.2002.120905> PMID: 11852419
- [4] Ra'awji BAA, Almogbel ES, Alharbi LA, Alotaibi AK, Al-Qazlan FA, Saquib J. Knowledge, attitudes, and practices of health-care workers regarding hand hygiene guidelines in Al-Qassim, Saudi Arabia: A multicenter study. *Int J Health Sci* 2018; 12(2): 3-8. PMID: 29599687
- [5] Hamid Ali H, Mustafa MM, Al-Rasheedi M, Balkhi B, Suliman N, Alshaafee W. Assessment of hospital staff knowledge, attitudes and practices (KAPS) on activities related to prevention and control of hospital acquired infections. *Int J Prev Treat* 2019; 8(1):

- 1-7.
<http://dx.doi.org/10.5923/j.ijpt.20190801.01>
- [6] Abalkhail A, Al Imam MH, Elmosaad YM, *et al.* Knowledge, attitude and practice of standard infection control precautions among health-care workers in a university hospital in Qassim, Saudi Arabia: A cross-sectional survey. *Int J Environ Res Public Health* 2021; 18(22): 11831.
<http://dx.doi.org/10.3390/ijerph182211831> PMID: 34831585
- [7] World Health Organization. Standard precautions for the prevention and control of infections: aide-memoire. 2022. Available from: <https://www.who.int/citations/i-item/WHO-UHL-IHS-IPC-2022.1>
- [8] Centers for Disease Control and Prevention. Core infection prevention and control practices for safe healthcare delivery in all settings. 2024. Available from: <https://www.cdc.gov/infection-control/hcp/core-practices/index.html>
- [9] Centers for Disease Control and Prevention. Standard precautions for all patient care. 2025. Available from: <https://www.cdc.gov/infection-control/hcp/basics/standard-precautions.html>
- [10] Alshammari SA, Alrasheed SS, Alruhaimi WA, Albnyan AI, Alruhaimi B, Hajj M. Knowledge, attitude, and practice of standard infection control precautions among medical students at King Khalid University Hospital. *Cureus* 2024; 16(6): e62768.
<http://dx.doi.org/10.7759/cureus.62768> PMID: 39036126
- [11] Yilma M, Taye G, Abebe W. Magnitude of standard precautions practices among healthcare workers in health facilities of Low and Middle Income Countries: A systematic review and meta-analysis. *PLoS One* 2024; 19(4): e0302282.
<http://dx.doi.org/10.1371/journal.pone.0302282> PMID: 38687766
- [12] Yezli S, Alotaibi B, Al-Abdely H, *et al.* Acquisition of respiratory and gastrointestinal pathogens among health care workers during the 2015 Hajj season. *Am J Infect Control* 2019; 47(9): 1071-6.
<http://dx.doi.org/10.1016/j.ajic.2019.02.033> PMID: 30979562
- [13] Alshamrani MM, Tannous E, Othman F, Al Zunitan M, Abalkhail M, El-Saed A. Competency level and determinants among infection prevention and control staff in the Middle East and North Africa region. *BMC Public Health* 2024; 24(1): 2224.
<http://dx.doi.org/10.1186/s12889-024-19717-x> PMID: 39148097
- [14] Alqahtani NS, Elmahboub RAA, Al-Qahtani FS, Al Qahtani SSAJ, Eldeeb SM, Al Margan AM. Assessment of healthcare workers' adherence to infection prevention and control measures in Najran City, Saudi Arabia, in the post-COVID-19 pandemic era. *J Pure Appl Microbiol* 2024; 18(2): 1235-45.
<http://dx.doi.org/10.22207/JPAM.18.2.43>
- [15] AlDakhil L, Yenugadhathi N, Al-Seraihi O, Al-Zoughool M. Prevalence and associated factors for needlestick and sharp injuries (NSIs) among dental assistants in Jeddah, Saudi Arabia. *Environ Health Prev Med* 2019; 24(1): 60.
<http://dx.doi.org/10.1186/s12199-019-0815-7> PMID: 31601166
- [16] Alanzi ME, Albalawi MAH, Kabrah S, Aljehani YT, Okashah AM, Aljohani ZDE. Knowledge, attitudes, and practices (KAPs) of healthcare workers towards MERS-CoV infection at PHCs in Madinah, KSA during Hajj 1440, 2019. *Am J Microbiol Res* 2019; 7(4): 122-9.
<http://dx.doi.org/10.12691/ajmr-7-4-4>
- [17] Tumala RB, Almazan J, Alabdulaziz H, *et al.* Assessment of nursing students' perceptions of their training hospital's infection prevention climate: A multi-university study in Saudi Arabia. *Nurse Educ Today* 2019; 81: 72-7.
<http://dx.doi.org/10.1016/j.nedt.2019.07.003> PMID: 31330405
- [18] Yosef T. Healthcare professionals' knowledge, attitude and practice of infection prevention in southwest Ethiopia. *Environ Health Insights* 2023; 17: 11786302231218819.
<http://dx.doi.org/10.1177/11786302231218819> PMID: 38116490
- [19] Sharma M, Bachani R. Knowledge, attitude, practice, and perceived barriers for the compliance of standard precautions among medical and nursing students in Central India. *Int J Environ Res Public Health* 2023; 20(8): 5487.
<http://dx.doi.org/10.3390/ijerph20085487> PMID: 37107771
- [20] Mohd-Nor N, Bit-Lian Y. Knowledge, attitude and practices of standard precaution among nurses in Middle-East hospital. *SciMedicine Journal* 2019; 1(4): 189-98.
<http://dx.doi.org/10.28991/SciMedJ-2019-0104-4>
- [21] Paul E, Alzaydani Asiri IA, Al-Hakami A, *et al.* Healthcare workers' perspectives on healthcare-associated infections and infection control practices: A video-reflexive ethnography study in the Asir region of Saudi Arabia. *Antimicrob Resist Infect Control* 2020; 9(1): 110.
<http://dx.doi.org/10.1186/s13756-020-00756-z> PMID: 32678049
- [22] Ni L, Yin X, Liu L, Sun K, Zhang W. Research on the effectiveness of the training of nosocomial infection control specialist nurses under the background of the new crown epidemic based on competence-based theory. *Altern Ther Health Med* 2024; 30: AT10298.
PMID: 38758138
- [23] Sharma R, Kumar P, Patel B, Gajjar S. Impact evaluation of domains of learning on universal work precautions among nursing staff in a tertiary care hospital, Western India. *Indian J Community Med* 2016; 41(2): 141-5.
<http://dx.doi.org/10.4103/0970-0218.153887> PMID: 27051089
- [24] Yabrodi M, Abulebda K, Pearson KJ, *et al.* Using a serious game to teach central line care in pediatric critical care nursing. *Am J Nurs* 2025; 125(2): 36-42.
<http://dx.doi.org/10.1097/AJN.0000000000000008> PMID: 39844230
- [25] Senbato FR, Wolde D, Belina M, *et al.* Compliance with infection prevention and control standard precautions and factors associated with noncompliance among healthcare workers working in public hospitals in Addis Ababa, Ethiopia. *Antimicrob Resist Infect Control* 2024; 13(1): 32.
<http://dx.doi.org/10.1186/s13756-024-01381-w> PMID: 38475931

DISCLAIMER: The above article has been published, as is, ahead-of-print, to provide early visibility but is not the final version. Major publication processes like copyediting, proofing, typesetting and further review are still to be done and may lead to changes in the final published version, if it is eventually published. All legal disclaimers that apply to the final published article also apply to this ahead-of-print version.